arXiv:hep-ex/0605015v1 5 May 2006

CP AND CPT TESTS WITH THE KLOE DETECTOR

THE KLOE COLLABORATION[a]
presented by MARIANNA TESTA
Dipartimento di Fisica, Università degli studi di Roma "La Sapienza" and INFN Sezione di Roma, P.le A.Moro 2, 00185 - Rome, Italy

Neutral kaons provide one of the most sensitive system to CP and CPT violation. Tests on CP, CPT and quantum mechanics have been performed at KLOE operating at the DAΦNE e^+e^- collider. Results on the quantum interference in the channel $\phi \to K_S K_L \to \pi^+\pi^-\pi^+\pi^-$, the measurement of the BR($K_L \to \pi^+\pi^-$) and the related CP violating parameter ϵ are presented. Using the Bell-Steinberger relation, CPT violating parameters have been also obtained.

1 Introduction

The KLOE detector operates at DAΦNE, an e^+e^- collider working at the center of mass energy $W \sim m_\phi \sim 1.02$ GeV. The ϕ mesons are produced essentially at rest and decay to $K_S K_L$ (K^+K^-) $\sim 34\%$ ($\sim 49\%$) of the times. The K mesons are produced in a pure $J^{PC} = 1^{--}$ coherent quantum state, so that observation of a K_S in an event signals (tags) the presence of a K_L and vice-versa: highly pure, almost monochromatic, back-to-back K_S and K_L beams can be obtained. Moreover K_S and K_L are distinguishable on the basis of their decay length: $\lambda_S \sim 0.6$ cm and $\lambda_L \sim 340$ cm.

The KLOE experiment is designed to exploit the unique feature of a ϕ-factory environment for the measurement of CP and CPT violation in the $K^0 - \bar{K}^0$ system and more generally for the study of kaons' decays and interference. The KLOE detector consists essentially of a drift chamber (DCH), surrounded by an electromagnetic calorimeter (EMC). A superconducting coil surrounding the barrel provides a 0.52 T magnetic field. Descriptions of the EMC and DCH can be found in ref. [1,2]

---

[a] F. Ambrosino, A. Antonelli, M. Antonelli, C. Bacci, P. Beltrame, G. Bencivenni, S. Bertolucci, C. Bini, C. Bloise, V. Bocci, F. Bossi, D. Bowring, P. Branchini, R. Caloi, P. Campana, G. Capon, T. Capussela, F. Ceradini, S. Chi, G. Chiefari, P. Ciambrone, S. Conetti, E. De Lucia, A. De Santis, P. De Simone, G. De Zorzi, S. Dell'Agnello, A. Denig, A. Di Domenico, C. Di Donato, S. Di Falco, B. Di Micco, A. Doria, M. Dreucci, G. Felici, A. Ferrari, M. L. Ferrer, G. Finocchiaro, S. Fiore, C. Forti, P. Franzini, C. Gatti, P. Gauzzi, S. Giovannella, E. Gorini, E. Graziani, M. Incagli, W. Kluge, V. Kulikov, F. Lacava, G. Lanfranchi, J. Lee-Franzini, D. Leone, M. Martini, P. Massarotti, W. Mei, S. Meola, S. Miscetti, M. Moulson, S. Müller, F. Murtas, M. Napolitano, F. Nguyen, M. Palutan, E. Pasqualucci, A. Passeri, V. Patera, F. Perfetto, L. Pontecorvo, M. Primavera, P. Santangelo, E. Santovetti, G. Saracino, B. Sciascia, A. Sciubba, F. Scuri, I. Sfiligoi, T. Spadaro, M. Testa, L. Tortora, P. Valente, B. Valeriani, G. Venanzoni, S. Veneziano, A. Ventura, R.Versaci, G. Xu

2 Quantum interference in the channel $K_L K_S \to \pi^+\pi^-\pi^+\pi^-$.

Test of quantum mechanics (QM) can be performed by studying the time evolution of the quantum correlated kaon system, in particular studying the interference pattern of the decay $K_L K_S \to \pi^+\pi^-\pi^+\pi^-$. According to QM, the distribution of the difference of the decay times $I(\Delta t)$ of the two kaons shows a characteristic destructive interference which prevents the two kaons from decaying into the same final state at the same time. As suggested in ref. [3,4], a simple way to parametrize a possible deviation of QM is to introduce a decoherence parameter $\zeta_{S,L}$ ($\zeta_{S,L} = 0$ in QM) as follows:

$$I(|\Delta t|) \propto e^{-|\Delta t|\Gamma_L} + e^{-|\Delta t|\Gamma_S} - 2\,\underbrace{(1 - \zeta_{S,L})}_{\text{decoherence}}\, cos(\Delta m|\Delta t|)e^{-\frac{\Gamma_s+\Gamma_L}{2}|\Delta t|} \tag{1}$$

Selecting a pure sample of $K_L K_S \to \pi^+\pi^-\pi^+\pi^-$ and fitting eq. 1 to data, KLOE has obtained the following preliminary result:



Fig. 1: Fit of the difference Δt of the decay times of $K_S \to \pi^+\pi^-$ and $K_L \to \pi^+\pi^-$. The black points are the data and the red ones are the results of the fit. The peak at $\Delta t \sim 17\tau_S$ is due to the regeneration on the beam pipe.

$$\zeta_{S,L} = 0.043\,^{+0.038}_{-0.035}{}_{\text{stat}} \pm 0.008_{\text{syst}},$$

consistent with QM predictions. The result of the fit is shown in fig. 1.

3 BR($K_L \to \pi^+\pi^-$)

KLOE has measured the BR($K_L \to \pi^+\pi^-$) using a K_L beam tagged by $K_S \to \pi^+\pi^-$ decays. The number of $K_L \to \pi^+\pi^-$ is obtained from a fit to the $\sqrt{E^2_{miss} + |\mathbf{p}_{miss}|^2}$ distribution, where



Fig. 2: $|\varepsilon|$ constraints by the measurements of $|V_{ub}|/|V_{uc}|$, Δm_d and by the limit on Δm_son the $(\bar{\rho}, \bar{\eta})$ from ref. [11], compared with the value of $|\varepsilon|$ from the measurements of BR($K_L \to \pi^+\pi^-$).

E_{miss} is the missing energy in the hypothesis of $K_L \to \pi^+\pi^-$ decay and $\mathbf{p}_{miss}$ is the missing momentum, with a linear combination of Monte Carlo distributions for $K_L \to \pi^+\pi^-$, $K_L \to \pi^\pm e^\mp \nu$, $K_L \to \pi^\pm \mu^\mp \nu$ $K_L \to \pi^+\pi^-\pi^0$ events, inclusive with respect to final-state radiation. The number of signal events has been normalized to the number of $K_L \to \pi\mu\nu$, in order to minimize systematic uncertainties on the tagging and tracking efficiency evaluation (exploiting the similar topology of the decays as well as the momentum overlap). Correcting for the tagging and tracking efficiency and using the BR($K_L \to \pi^\pm\mu^\mp\nu$) from ref. [7], we obtain: BR($K_L \to \pi^+\pi^-$) = $(1.963 \pm 0.012_{\text{stat}} \pm 0.017_{\text{syst}}) \times 10^{-3}$. The result is in good agreement with the measurement of KTeV [5] $(1.975 \pm 0.012) \times 10^{-3}$ and in strong disagreement with that reported by the PDG [6], $(2.090 \pm 0.025) \times 10^3$. This result can be used to determine $|\eta_{+-}|$ and $|\varepsilon|$ correcting for the small contribution of ε'. Using the measurements of BR($K_S \to \pi^+\pi^-$) and τ_{K_L} from KLOE [9,7,8], and the value of τ_{K_S} from PDG[6] and subtracting the contribution of the photon direct emission [10] to the value of BR($K_L \to \pi^+\pi^-$), we obtain: $|\eta_{+-}| = (2.219 \pm 0.013) \times 10^{-3}$. Finally, using the world average measurement of $\text{Re}(\varepsilon'/\varepsilon) = (1.67 \pm 0.26) \times 10^{-3}$, and assuming equal phases between ε' and ε we obtain $|\varepsilon| = (2.216 \pm 0.013) \times 10^{-3}$, in disagreement with the value $|\varepsilon| = (2.284 \pm 0.014) \times 10^{-3}$ reported in ref. [6]. The value of $|\varepsilon|$ can be can be compared with the prediction [11] $|\varepsilon| = (2.875 \pm 0.455) \times 10^{-3}$ where to test the mechanism of the CP violation in the Standard Model, the value of $|\varepsilon|$ has been computed from the measurement of the CP conserving observables: Δm_d, Δm_s, V_{ub}, and V_{cb}. No significant deviation from the Standard Model prediction has been observed.

4 Bell Steinberger Relation

The most powerful test of CPT invariance in the neutral kaon system is presently obtained by means of the Bell-Steinberger relation [12], which relates CPT and CP violating parameters, $Im(\delta)$ and $Re(\epsilon)$, to the decay amplitudes of K_L and K_S into the same final state:

$$(1 + i \tan \phi_{SW})[\text{Re}(\epsilon) - i\,\text{Im}(\delta)] = \sum_{\substack{\text{final}\\ \text{states } f}} A(K_L \to f)^\star A(K_S \to f)/\Gamma_S = \sum_{\substack{\text{final}\\ \text{states } f}} \alpha_f \qquad (2)$$

where ϕ_{SW} is the superweak phase, defined by $\tan \phi_{SW} = 2\Delta M/(\Gamma_S - \Gamma_L)$. For the determination of the α_f parameters, experimental inputs are K_L and K_S branching ratios, the relative phases between the amplitudes, and the K_L and K_S lifetimes, τ_{K_S} and τ_{K_L}. We use the value of τ_{K_S} reported by the PDG [6] and τ_{K_L} from KLOE average [7,8] and the following measurements:

- the new KLOE measurement of BR($K_S \rightarrow \pi^+\pi^-$), BR($K_S \rightarrow \pi^0\pi^0$) from ref [9] which enters in the evaluation of $|\alpha_{+-}|$ and $|\alpha_{00}|$,
- the average between the BR($K_L \rightarrow \pi^+\pi^-$) here presented and that measured by KTeV[5], used to determine $|\alpha_{+-}|$,
- the measurement of BR($K_L \rightarrow \pi^0\pi^0$) from KTeV [5], used to determine $|\alpha_{00}|$,
- the values, ϕ_{+-} and ϕ_{00}, of the phases of α_{+-} and α_{00}, taken from the PDG[6] fit without assuming CPT symmetry,
- the measurement of the CP conserving direct component contribution to the process $K_L \rightarrow \pi^+\pi^-\gamma$ from ref [10] and the upper limit on the direct component contribution to the process $K_S \rightarrow \pi^+\pi^-\gamma$ [6], both entering in the evaluation of $\alpha_{+-\gamma}$,
- the recent KLOE upper limit on the BR($K_S \rightarrow \pi^0\pi^0\pi^0$)[13], which constraints the value of $|\alpha_{000}|$,
- the measurement of the BR($K_S \rightarrow \pi^+\pi^-\pi^0$) reported in the PDG [6],
- the recent KLOE measurement of the semileptonic K_S charge asymmetry A_S [9], which allows to calculate the semileptonic contribution $\alpha_{kl3} = 2\tau_{K_S}/\tau_{K_L} B(kl3)((A_S + A_L)/4 - iIm(\delta) + Im(x+))$, being $x+$ the parameter describing the $\Delta S = \Delta Q$ violation in the semileptonic decays. $Im(x+)$ has been determined from a combined fit of A_S with the semileptonic time dependent decay rate asymmetry measured by CPLEAR [14]. The semileptonic K_L charge asymmetry A_L has been taken from the PDG [6].

As $\phi_{+-\gamma}$, the phase of $\alpha_{+-\gamma}$, has not been measured yet, no constraints have been assumed on its value. Using these experimental inputs, the differences between the K_S and K_L masses and lifetimes, ΔM and $\Delta\Gamma$, reported in the PDG[6] (for the determination of ϕ_{SW}), we obtain: Re(ϵ) = $(160.2 \pm 1.3) \times 10^{-5}$ and Im(δ) = $(1.2 \pm 1.9) \times 10^{-5}$, resulting in a considerable improvement to the CPLEAR measurement [15]: Re(ϵ) = $(164.9 \pm 2.5) \times 10^{-5}$ and Im(δ) = $(2.4 \pm 5.0) \times 10^{-5}$.

References

1. M. Adinolfi *et al.*, KLOE Collaboration, *Nucl. Instrum. Methods* A **488**, 51 (2002)
2. M. Adinolfi *et al.*, KLOE Collaboration, *Nucl. Instrum. Methods* A **482**, 364 (2002)
3. R. A. Bertlmann *et al.*, *Phys. Rev.* D **60**, 114032 (1999)
4. P. H. Eberhard, *Test of quantum mechanics at a ϕ factory*, vol.1. The Second DAΦNE Physics Handbook; eds. L. Maiani *et. al.*, (SIS-Publ. dei Laboratori Nazionali di Frascati), Frascati , Italy (1995)
5. KTeV Collaboration, T.Alexopolous *et al.*, *Phys. Rev.* D **70**, 0920006 (2004)
6. Particle Data Group, S. Eidelman *et al.*, *Phys. Lett.* B **592**, 1 (2004)
7. KLOE Collaboration, F. Ambrosino *et al.*, *Phys. Lett.* B **632**, 43 (2006)
8. KLOE Collaboration, F. Ambrosino *et al.*, *Phys. Lett.* B **626**, 15 (2006)
9. KLOE Collaboration, F. Ambrosino *et al.*, Precise measurement of $\Gamma(K_S \rightarrow \pi^+\pi^-)/\Gamma(K_S \rightarrow \pi^0\pi^0)$ with the KLOE detector at DAΦNE hep-ex/0601025
10. E. Ramberg *et al.*, *Phys. Rev. Lett.* **70**, 2525 (1993)
11. UTfit Collaboration (http://www.utfit.org), private communication
12. J.S. Bell and J. Steinberger, Proc. of Oxford Int. Conf. on Elementary Part., 195 (1965)
13. KLOE collaboration, F. Ambrosino *et al.*, *Phys. Lett.* B **619**, 61 (2005)
14. CPLEAR Collaboration, Angelopoulos *et al.*, EPJ **22**, 55 (2001)
15. CPLEAR Collaboration, Apostolakis *et al.*, *Phys. Lett.* B **456**, 297 (1999)